

- Flack, J. M., Victor, R., Watson, K., Ferdinand, K. C., Saunders, E., Tarasenko, L., & Bruschi, P. (2008, January). Improved attainment of blood pressure and cholesterol goals using single-pill amlodipine/atorvastatin in African Americans: the CAPABLE trial. In Mayo Clinic Proceedings (Vol. 83, No. 1, pp. 35-45). Elsevier.
- Fox, C. S., Pencina, M. J., Wilson, P. W., Paynter, N. P., Vasan, R. S., & D'Agostino, R. B. (2008). Lifetime risk of cardiovascular disease among individuals with and without diabetes stratified by obesity status in the Framingham heart study. *Diabetes care*, 31(8), 1582-1584.
- Fujimoto, W. Y., Bergstrom, R. W., Leonetti, D. L., Newell-Morris, L. L., Shuman, W. P., & Wahl, P. W. (1994). Metabolic and adipose risk factors for NIDDM and coronary disease in third-generation Japanese-American men and women with impaired glucose tolerance. *Diabetologia*, 37(5), 524-532.
- Gæde, P., Lund-Andersen, H., Parving, H. H., & Pedersen, O. (2008). Effect of a multifactorial intervention on mortality in type 2 diabetes. *New England journal of medicine*, 358(6), 580-591.
- Ginsberg, H. N., Elam, M. B., Lovato, L. C., Crouse 3rd, J. R., Leiter, L. A., Linz, P., & Wright, D. (2010). Effects of combination lipid therapy in type 2 diabetes mellitus. *The New England journal of medicine*, 362(17), 1563-1574.
- Gottschalk, M., Danne, T., Vlajnic, A., & Cara, J. F. (2007). Glimepiride Versus Metformin as Monotherapy in Pediatric Patients With Type 2 Diabetes A randomized, single-blind comparative study. *Diabetes Care*, 30(4), 790-794.