

Nava-Ocampo AA, Velázquez-Armenta EY, Han JY, Koren G. Use of proton pump inhibitors during pregnancy and breastfeeding. Can Fam Physician. 2006; 52: 853-4 .

Nichols DJ, Muirhead GJ, Harness JA. Pharmacokinetics of sildenafil after single oral doses in healthy male subjects: absolute bioavailability, food effects and dose proportionality. Br J Clin Pharmacol 2002; 53 (Suppl. 1): 5S-12S.

Onal A.; Oztunç A. Development and validation of high performance liquid chromatographic method for the determination of esomeprazole in tablets, J. Food Drug Anal., Vol. 14, No. 1, 2006, pp 12-18.

Patel, B.H.; Suhagia, B.N.; Patel, M.M.; Patel, J.R. Determination of pantoprazole, domperidone, rabeprazole, esomeprazole and itopride in pharmaceutical products by reverse phase liquid chromatography using single mobile phase, Chromatographia, 65(2007)11-12, pp-743-748.

Patil Shamkant S, Dhabale Pandurang N and Kuchekar Bhanudas S. Development and Statistical Validation of Spectrophotometric Method for Estimation of Esomeprazole in Tablet Dosage Form. Asian J. ResearchChem. 2(2): April.-June, 2009.

Patterson B, Bedding A, Jewell H et al. The effect of intrinsic and extrinsic factors on the pharmacokinetic properties of tadalafil (IC351). Int J Impot Res 2001;.