

(GMDP) and enhancement of absorption in rat by delivery in a water-in-oil microemulsion. *Int J Pharm*, 199(1), 17-28.

- Ma, Z., Lim, T.M., & Lim, L.Y. (2005). Pharmacological activity of peroral chitosan-insulin nanoparticles in diabetic rats. *Int.J.Pharm*, 293(2), 271-280.
- Makhlof, A., Tozuka, Y., & Takeuchi, H. (2011). Design and evaluation of novel pH-sensitive chitosan nanoparticles for oral insulin delivery. *Eur J Pharm Sci.*, 42(5), 445-451.
- Marguerite, R. (2006). Chitin and Chitosan: Properties and application. *Progress in Polymer Science*, 31(7), 603-632.
- Marshal, S. (1985). Kinetics of Insulin Receptor Internalization and Recycling in Adipocytes. *THE JOURNAL OF BIOLOGICAL CHEMISTRY*, 260(7), 4136-4144.
- Martinez, M., & Amidon, G. (2002). A Mechanistic Approach to Understanding the Factors Affecting Drug Absorption: A Review of Fundamentals. *The Journal Of Clinical Pharmacology*, 42(6), 620-643.
- Martins, S., Sarmento, B., Ferreira, D.C., & Souto, E.B. (2007). Lipid-based colloidal carriers for peptide and protein delivery – liposomes versus lipid nanoparticles. *International Journal of Nanomedicine*, 2(4), 595–607.
- Matsuzawa, A., Morishita, M., Takayama, K., & Nagai, T. (1995). Absorption of Insulin Using Water-in-Oil-in-Water Emulsion from an Enteral Loop in Rats. *Biological & Pharmaceutical Bulletin*, 18(12), 1718-1723.
- Meibohm, B. (2006). *Pharmacokinetics and pharmacodynamics of biotech drugs*. Weinheim: WILEY-VCH.
- Meier, J.J., Holst, J.J., Schmidt, W.E., & Nauck, M.A. (2007). Reduction of hepatic insulin clearance after oral glucose ingestion is not mediated by