

- Nakamura, K., Morishita, M., Ehara, J., Onuki, Y., Yamagata, T., Kamei, N., Lowman, A.M., Peppas, N.A., & Takayama K. (2008). Key functions in polymer carriers for intestinal absorption of insulin. *Int.J.Pharm*, 354(2), 135-142.
- Nakamura, K., Murray, R.J., Joseph. J.I., Peppas, N.A., Morishita, M., & Lowman, A.M. (2004). Oral insulin delivery using p(MAA-g-EG) hydrogels: effects of network morphology on insulin delivery characteristics. *J.Control Release*, 95(3), 589-599.
- Nayak, A.K. (2014). Advances in therapeutic protein production and delivery. *International Journal of Pharmacy and Pharmaceutical Sciences*, 2(2), 1-5.
- Ohno, T., Horio, F., Tanaka, S., Terada, M., Namikawa, T., & Kitch, J. (2000). Fatty liver and hyperlipidemia in IDDM (insulin dependent diabetes mellitus) of Streptozotocin treated shrews. *Life Sci.*, 66(2):125-31.
- Owens, D. (2002). New horizons — alternative routes for insulin therapy. *Nature Reviews Drug Discovery*, 1(7), 529-540.
- Owens, D.R., Vora, J.P., Birtwell, J., Luzio, S., & Hayas, T.M. (1988). The influence of aprotinin on regional absorption of soluble human insulin. *Br.J.Clin.Pharmacol*, 25(4), 453-456.
- Ozsoy, Y., Gungor, S., & Cevher, E. (2009). Nasal Delivery of High Molecular Weight Drugs. *Molecules*, 14(9), 3754-3779.
- Patel, H.M., Stevenson, R.W., Parons, J.A., & Ryman, B.E. (1982). Use of liposomes to aid intestinal absorption of entrapped insulin in normal and diabetic dogs. *Biochem. Biophys. Acta*, 716(2), 188-193.
- Patel, V., Liu, F., & Brown, M. (2011). Advances in oral transmucosal drug delivery. *Journal Of Controlled Release*, 153(2), 106-116.