

| | |
|--|-----|
| 3.2.4.5 Calibration curve 5 | 103 |
| 3.2.4.6 Calibration curve 6 | 104 |
| 3.2.5 Recovery | 107 |
| 3.2.5.1 Recovery in mobile phase | 107 |
| 3.2.5.2 Recovery in the serum | 107 |
| 3.3 Representative HPLC chromatograms of analyzed Sprague-Dawley rat serum and Krebs buffer samples | 110 |
| 3.4 <i>In vivo</i> experiments | 112 |
| 3.4.1 Effect of GlcN on PRN BA | 112 |
| 3.4.2 Effect of cimetidine and rifampin on PRN BA | 113 |
| 3.4.3 Effect of GlcN on PRN effective intestinal permeability (P_{eff}) and clearance | 115 |
| 3.5 Effect of GlcN on PRN <i>in situ</i> single-pass intestinal perfusion | 117 |
| 3.6 Effect of GlcN on PRN absorption in everted rat intestine sac (ERIS) | 119 |
| 3.7 Hepatocyte cell isolation and culture | 120 |
| 3.7.1 Effect of GlcN on PRN metabolism | 120 |
| 3.7.2 Effect of cimetidine and rifampin on PRN metabolism | 121 |
| Chapter Four | 122 |
| 4. Discussion | |
| Chapter Five | 130 |
| 5. Conclusion and future work | |