

calibration curve used in the calculation of measured concentration of intra-day one is shown in table 3.36 .	
Table 3.26 Intra-day accuracy & precision data for all QC samples LLOQ, QC low, QC mid, and QC high of propranolol based on the standard calibration curves of the second day of validation (n=6). Corresponding calibration curve used in the calculation of measured concentration of intra-day two is shown in table 3.37 .	90
Table 3.27 Intra-day accuracy & precision data for all QC samples LLOQ, QC low, QC mid, and QC high of propranolol based on the standard calibration curves of the third day of validation (n=6). Corresponding calibration curve used in the calculation of measured concentration of intra-day three is shown in table 3.38 .	91
Table 3.28 Inter-day accuracy & precision data for the three days of QC samples validation (LLOQ, QC low, QC mid, and QC high) of propranolol based on the standard calibration curves for the three days of validation (n=6).	93
Table 3.29 Propranolol QC low samples (150 ng/ml) results for freeze-thaw stability test (n=3).	94
Table 3.30 Propranolol QC high samples (2500 ng/ml) results for freeze-thaw stability test (n=3).	95
Table 3.31 Propranolol QC low samples (150 ng/ml) results for bench-top stability test (n=3).	96
Table 3.32 Propranolol QC high samples (2500 ng/ml) results for bench-top stability test (n=3).	96
Table 3.33 Propranolol QC low samples (150 ng/ml) results for autosampler stability test (n=3).	97
Table 3.34 Propranolol QC high samples (2500 ng/ml) results for autosampler stability test (n=3).	97
Table 3.35 Summary of all R ² , slope and intercept data of the six calibration curves.	98